



Value Life

Operating Room  
Tissue Adhesive

**INDERMIL<sup>®</sup> flexifuze<sup>™</sup>**

The next generation  
tissue adhesive



# INDERMIL<sup>®</sup> flexifuze<sup>™</sup>

## The next generation tissue adhesive

The ready-to-use, octyl blend cyanoacrylate product can be used for the closure of minor/low tension skin wounds, including surgical and laproscopic incisions, trauma induced lacerations and where appropriate can be combined with deep dermal stitches.

This rapid-setting adhesive seals the skin in just 30 seconds\* and will stay in place for five – eight days, whilst protecting the patient from microbial penetration and the risk of infection.

## Features and Benefits

● A **gentle** alternative for the patient

- **Reduced patient anxiety:** No need for a return visit to remove sutures or staples
- **Gentle on the skin:** No invasive skin punctures
- **Shower-proof:** Patients can shower as soon as adhesive has dried
- **0.75g ampoule:** Allows closure of multiple wounds on the same patient
- **No discernable heat during application:** No burning or stinging.



● An **easy** alternative for the clinician

- **No need to refrigerate:** Adhesive can be stored at room temperature (4-30°C)
- **Easy to use:** Includes flow control applicator with patented brush tip
- Adhesive has a **long working life** (up to 20 minutes)
- Seals wound **against microbial penetration**
- May eliminate the use of sharps and therefore **reduces the risk of needle-stick injury.**

\*30 seconds is a benchtop standard test used by TA manufacturers. Allow for 30 – 60 seconds drying time to ensure the adhesive is tack-free.



# Instructions for Use

For additional product information, warnings, precautions and contraindications, please refer to the 'Instructions for Use' leaflets supplied with the product.

## 1. Wound preparation

Remove any foreign bodies if present. Clean and dry the wound thoroughly, ensuring wound edges are easily apposed.

## 2. Opening the product

Remember the contents of the blister pack are sterile prior to opening. Slowly peel open blister pack with label facing upwards to expose the device.

## 3. Application to the wound

If desired attach the brush applicator onto the device.

Hold the device in one hand with the tip pointing upwards away from the patient and user to avoid inadvertent contact. Gently pinch the device towards the centre of the tube to crush the inner glass ampoule to release the adhesive.

When using the brush tip, ensure that the centre of the brush is directly above the wound.

## 4. The closed wound

Hold wound edges together with fingers, forceps or other appropriate methods. Apply adhesive sparingly along the length of the wound site. Avoid adhesive entering the wound. Continue holding wound edges together to allow adhesive to set for approximately 30 seconds. Wounds that are subject to tension can be secured by subcuticular sutures prior to application of adhesive.

INDERMIL® flexifuze™ tissue adhesive dries to form a strong, colourless seal within seconds.

## 5. Protecting the wound


Protective dry dressing such as gauze can be applied to the wound once adhesive is completely dry.

## 6. Disposal

Dispose of ampoule safely in sharps bin or clinical waste.

## 7. The result

The healing result will depend on individual circumstances relevant to the specific wound being closed.




Please visit [www.ndermilflexifuze.co.uk](http://www.ndermilflexifuze.co.uk) to view our how to use INDERMIL flexifuze video.



Patient advice sheet  
Code: DXJB0006001



Bravery stickers  
Code: DXJP0006003



Brave baby certificate  
Code: DXJP0006001



Brave child certificate  
Code: DXJP0006002

# Product Support

Vygon is committed to education and support for our customers. We provide in-service training sessions for all staff and all shift patterns, to help ensure your clinicians achieve best practice when

delivering care to patients. In addition, patient advice leaflets and children's bravery certificates are available on request, to help assist the clinicians in supporting and informing their patients.

# Order INDERMIL<sup>®</sup> flexifuze<sup>™</sup> today

Product Codes		Description	Box Quantity
Vygon	NHSSC		
CM001	FVT150	INDERMIL <sup>®</sup> flexifuze <sup>™</sup> Tissue Adhesive	12

## Technical Support Data for tissue adhesive

1. Roberts AC. The Tissue Adhesive INDERMIL<sup>®</sup> and its use in Surgery. *Acta Chirurgiae Plasticae* 1998, 40 (1): 22-25
2. Sinha S, Naik M, Wright V, Timmons J, Campbell AC. A Single Blind, Prospective, Randomised Trial Comparing n-butyl 2-cyanoacrylate Tissue Adhesive (INDERMIL<sup>®</sup>) and Sutures for Skin Closure in Hand Surgery. *Journal of Hand Surgery (Br)* 2001, 26 (3): 264-5
3. Schonauer F, Pereira J, La Rusca I, Harris J, Cullen K. Use of INDERMIL<sup>®</sup> Tissue Adhesive for Closure of Superficial Skin Lacerations in Children. *Minerva Chirurgica* 2001, 56: 427-9
4. Eckersley E. Treating the Child with Minor Scalp Wounds. *Emergency Nurse* 1998, 6, No.5
5. Doraiswamy NV, Baig H, Hammett S, Hutton M. Which Tissue Adhesive for Wounds? *Injury* 2003, 34 (8): 564-7
6. Jaibaji M, Liddington M, Geary P, Darcy C, Batchelor AG, Roberts AC. Evaluation of Long-Term Outcome of Wounds Closed with INDERMIL<sup>®</sup> Adhesive. *Eur J Plast Surg* 2000, 23: 330-2
7. Qureshi A, Roberts AC, Monson JRT, Drew PJ, Duthie GS. N-butyl cyanoacrylate Adhesive for Skin Closure of Abdominal Wounds: Preliminary Results. *Ann R Coll Surg Engl* 1997, 79: 414-5
8. Rogerson L, Mason GC, Roberts AC. Preliminary Experience with Twenty Perineal Repairs using INDERMIL<sup>®</sup> Tissue Adhesive. *Eur J Obstet Gynecol and Reprod Biol* 2000, 88: 139-42
9. Samuel PR, Roberts AC, Nigam A. The Use of INDERMIL<sup>®</sup> (n-butyl cyanoacrylate) in Otorhinolaryngology and Head and Neck Surgery. A Preliminary Report on the First 33 Patients. *Journal of Laryngology and Otology* 1997, III: 536-40
10. England RJ, Roberts AC, Raines CH. Does INDERMIL<sup>®</sup> Glue Improve Success Rates in Myringoplasty? Interim Analysis of a Prospective Trial. *Rev. Laryngol Otol Rhinol Bord* 2000, 121 (2): 91-3



This literature has been made using carbon balanced paper. If you would like to find out more about what Vygon (UK) Ltd does for the environment, please visit: [www.vygon.co.uk/company/corporate](http://www.vygon.co.uk/company/corporate)

**Manufactured by Connexicon Medical**

**For further information, please contact: [vygon@vygon.co.uk](mailto:vygon@vygon.co.uk)**

The specifications shown in this leaflet are for information only and are not, under any circumstances, of a contractual nature. This document is intended for use in the UK only

Vygon (UK) Ltd • The Pierre Simonet Building • V Park • Gateway North  
• Latham Road • Swindon • Wiltshire • SN25 4DL  
Tel: 01793 748800 • Fax: 01793 748899 • Twitter: @vygonuk  
Web: [www.vygon.co.uk](http://www.vygon.co.uk) Content correct as of 01/2017 Code: DXJB0006002 v2  
Copyright Vygon (UK) Ltd 2017

**connexicon**  
medical



[www.vygon.co.uk](http://www.vygon.co.uk)