

# Intravesical Instillation


## Context

A local treatment by vesical instillation of Mitomycine MMC (cytotoxic drug) or BCG (bacillus Calmette-Guérin) is often recommended after Transurethral Resection of Bladder Tumor (TURBT). However handling these toxic drugs requires a high level of security to protect healthcare workers and their environment. Recognised risks are Chemical and bacteriological contamination.



## Problem

Drug leakage and/or splashing during preparation, transfer, instillation and disposal.



## Solution: qimocap - qimo♂ - qimoadapt

For the handling of toxic drugs, the Qimono range ensures a high level of security for the healthcare workers from the drug vial to the disposal of the urinary catheter.



**qimocap 20mm**  
Ordering code: 7219.02



**qimo♂**  
Ordering code: 7219.91



**qimoadapt**  
Ordering code: 7210.06

## INTRAVASCULAR THERAPIES

Vygon (UK) Ltd, The Pierre Simonet Building, V Park, Gateway North, Latham Road, Swindon, Wiltshire SN25 4DL

Tel: 01793 748800 Fax: 01793 748899 Email: [vygon@vygon.co.uk](mailto:vygon@vygon.co.uk)

[vygon.co.uk](http://vygon.co.uk) [@vygonuk](https://twitter.com/vygonuk) [vygonuk](https://www.linkedin.com/company/vygonuk) [vygonuk](https://www.youtube.com/channel/UCvYgONk) [vygonuk](https://www.instagram.com/vygonuk)

Copyright Vygon (UK) Ltd 2020 Content correct as of: 03/2020 Code: V00678 v1



Value Life